

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification 7 : C07D 487/04, A61K 31/415 // (C07D 487/04, 235:00, 221:00) (C07D 487/04, 235:00, 223:00)</p>	<p>A1</p>	<p>(11) International Publication Number: WO 00/69863 (43) International Publication Date: 23 November 2000 (23.11.00)</p>
<p>(21) International Application Number: PCT/US00/13413 (22) International Filing Date: 16 May 2000 (16.05.00) (30) Priority Data: 60/134,822 19 May 1999 (19.05.99) US (71) Applicant (for all designated States except US): THE PROCTER & GAMBLE COMPANY [US/US]; One Procter & Gamble Plaza, Cincinnati, OH 45202 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): LIU, Song [CN/US]; 9338 Ambleside Drive, Cincinnati, OH 45241 (US). PORTLOCK, David, Edward [US/US]; 3237 Chestnut Landing Drive, Maineville, OH 45039 (US). PONG, Schwe, Fang [US/US]; 8005 Tylers Way, West Chester, OH 45069 (US). (74) Agents: REED, T., David et al.; The Procter & Gamble Company, 5299 Spring Grove Avenue, Cincinnati, OH 45217-1087 (US).</p>	<p>(81) Designated States: AE, AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), DM, EE, EE (Utility model), ES, FI, FI (Utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	
<p>(54) Title: IMIDAZO-CONTAINING HETEROCYCLIC COMPOUNDS, THEIR COMPOSITIONS AND USES</p> <div data-bbox="462 1144 1063 1333" data-label="Chemical-Block"> </div> <p>(57) Abstract</p> <p>The subject invention involves compounds having structure (I), wherein each R1 is independently alkyl, aryl, or heterocycle; each R2 - R7 is independently hydrogen or other substituent; B is nil and n is 0-3, or B is ethenyl and n is 0-1; and pharmaceutically acceptable forms thereof. The subject invention also involves pharmaceutical compositions containing such compounds, and methods for treating or preventing diseases or disorders using such compounds.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

IMIDAZO-CONTAINING HETEROCYCLIC COMPOUNDS, THEIR COMPOSITIONS AND USES

FIELD OF THE INVENTION


The subject invention relates to novel imidazo-containing heterocyclic compounds, pharmaceutical compositions containing them, and their therapeutic or preventative use in the areas of cardiovascular, oncology, infectious and inflammatory diseases.

BACKGROUND

Certain imidazo-isoquinoline compounds are disclosed in U.S. Patent No. 3,917,610 issued November 4, 1975; they are reported to have certain cardiovascular activities. One such compound is further reported on in Borchard, Fox, and Greeff, "The Positive Inotropic, Antiarrhythmic and Na⁺, K⁺-ATPase Inhibitory Effects of the Isoquinoline Derivative, BILA", *Archives of Pharmacology*, vol. 312 (1980), pp. 187-192.

SUMMARY OF THE INVENTION

The subject invention includes compounds having the structure:



wherein:

- (a) each R1 is independently selected from alkyl, aryl, and heterocycle;
- (b) R2 is selected from hydrogen, alkyl, alkylacyl, arylacyl, alkylsulfonyl and arylsulfonyl;

- (c) each R3 is independently selected from hydrogen, halo, alkyl, aryl, heterocycle, nitro, cyano, and unsubstituted or alkyl- or aryl- or heterocycle-substituted hydroxy, thio, amino, amide, formyl (acyl), carboxy, and carboxamide; two R3's on adjacent carbons may optionally together be alkylene or heteroalkylene, thereby forming a fused ring with the phenyl to which they are attached;
- (d) R4 is selected from hydrogen, halo, alkyl, aryl, heterocycle, and carboxy and its alkyl and aryl esters and amides;
- (e) each R5 is independently selected from hydrogen, alkyl, and aryl;
- (f) each R6 is independently selected from hydrogen, halo, alkyl, aryl, heterocycle, nitro, cyano, and unsubstituted or alkyl- or aryl- or heterocycle-substituted hydroxy, thio, amino, amide, sulfonamide, formyl (acyl), carboxy, and carboxamide; or the two R6's may optionally together be alkylene or heteroalkylene, thereby forming a fused ring with the phenyl to which they are attached;
- (g) B is nil or $-C(R7)=C(R7)-$;
- (h) if B is nil, n is an integer from 0 to about 3, otherwise n is from 0 to about 1;
- (j) each R7 is independently selected from hydrogen, alkyl and aryl;

and an optical isomer, diastereomer or enantiomer thereof; a pharmaceutically acceptable salt, hydrate, or biohydrolyzable ester, amide or imide thereof.

The subject invention also includes compositions comprising a subject compound and a pharmaceutically-acceptable excipient; and methods for treating or preventing diseases or disorders by administering to a human or lower animal in need thereof, a safe and effective amount of a subject compound.

DETAILED DESCRIPTION OF THE INVENTION

As used herein unless specified otherwise, "alkyl" means a hydrocarbon chain which is branched, linear or cyclic, saturated or unsaturated (but not aromatic), substituted or unsubstituted. The term "alkyl" may be used alone or as part of another word where it may be shortened to "alk" (e.g., in alkoxy, alkylacyl). Preferred linear alkyl

have from one to about twenty carbon atoms, more preferably from one to about ten carbon atoms, more preferably still from one to about six carbon atoms, still more preferably from one to about four carbon atoms; most preferred are methyl or ethyl. Preferred cyclic and branched alkyl have from three to about twenty carbon atoms, more preferably from three to about ten carbon atoms, more preferably still from three to about seven carbon atoms, still more preferably from three to about five carbon atoms. Preferred cyclic alkyl have one hydrocarbon ring, but may have two, three, or more, fused hydrocarbon rings. Preferred alkyl are unsaturated with from one to about three double or triple bonds, preferably double bonds; more preferably they are mono-unsaturated with one double bond. Still more preferred alkyl are saturated. Saturated alkyl are referred to herein as "alkanyl". Alkyl unsaturated only with one or more double bonds (no triple bonds) are referred to herein as "alkenyl". Preferred substituents of alkyl include halo, alkyl, aryl, heterocycle, hydroxy, alkoxy, aryloxy, thio, alkylthio, arylthio, amino, alkylamino, arylamino, amide, alkylamide, arylamide, formyl, alkylacyl, arylacyl, carboxy and its alkyl and aryl esters and amides, nitro, and cyano. Also, unsubstituted alkyl are preferred.

As used herein, "heteroatom" means a nitrogen, oxygen, or sulfur atom.


As used herein, "alkylene" means an alkyl which connects two other moieties, "heteroalkylene" means an alkylene having one or more heteroatoms in the connecting chain.

As used herein unless specified otherwise, "aryl" means an aromatic hydrocarbon ring (or fused rings) which is substituted or unsubstituted. The term "aryl" may be used alone or as part of another word (e.g., in aryloxy, arylacyl). Preferred aryl have from six to about fourteen, preferably to about ten, carbon atoms in the aromatic ring(s), and a total of from about six to about twenty, preferably to about twelve, carbon atoms. Preferred aryl is phenyl or naphthyl; most preferred is phenyl. Preferred substituents of aryl include halo, alkyl, aryl, heterocycle, hydroxy, alkoxy, aryloxy, thio, alkylthio, arylthio, amino, alkylamino, arylamino, amide, alkylamide, arylamide, formyl, alkylacyl, arylacyl, carboxy and its alkyl and aryl esters and amides, nitro, and cyano. Also, unsubstituted aryl are preferred.

As used herein unless specified otherwise, "heterocycle" means a saturated, unsaturated or aromatic cyclic hydrocarbon ring (or fused rings) with one or more heteroatoms in the hydrocarbon ring(s). Preferred heterocycles have from one to about six heteroatoms in the ring(s), more preferably one or two or three heteroatoms in the ring(s). Preferred heterocycles have from three to about fourteen, preferably to about ten, carbon plus heteroatoms in the ring(s), more preferably from three to about seven, more preferably still five or six, carbon plus heteroatoms in the rings(s); and a total of from three to about twenty carbon plus heteroatoms, more preferably from three to about ten, more preferably still five or six, carbon plus heteroatoms. Preferred heterocycles have one ring, but may have two, three, or more, fused rings. More preferred heterocycle rings include those which are one ring with 5 or 6 carbon plus heteroatoms in the ring with no more than three ring heteroatoms, no more than two of which are O and S. Still more preferred are such 5- or 6-ring atom heterocycles with one or two ring atoms being O or S and the others being C; or with one, two or three ring atoms being N and the others being C. Such preferred 5- or 6-ring atom heterocycles are preferably saturated, unsaturated with one or two double bonds, or aromatic. Such preferred 5- or 6-ring atom heterocycles are preferably a single ring; or fused with a 3- to 6-ring atom hydrocarbon ring which is saturated, unsaturated with one double bond, or aromatic (phenyl); or fused with another such 5- or 6-ring atom heterocyclic ring. Heterocycles are unsubstituted or substituted. Preferred heterocycle substituents are the same as for alkyl.

Compounds of the Invention

The subject invention involves compounds having the following structure:



In structure 1, each R1 is independently selected from alkyl, aryl, and heterocycle. Preferred R1 include linear alkanyl having from 1 to about 6 carbon atoms, linear alkenyl having from 2 to about 6 carbon atoms, and branched and cyclic alkanyl and alkenyl having from 3 to about 6 carbon atoms, such alkenyl preferably having 1 double bond. Such preferred alkanyl and alkenyl are preferably unsubstituted, or substituted with phenyl, heterocycle having 5 or 6 ring atoms, carboxy and its C₁ - C₆ alkyl and phenyl esters, or cyano. More preferably such alkanyl and alkenyl have up to about 5 carbon atoms, more preferably still up to 4 carbon atoms. More preferred R1 are methyl, ethyl and isopropyl. Most preferred R1 is unsubstituted methyl. Also preferred is both R1 being the same moiety.

In structure 1, R2 is selected from hydrogen, alkyl, alkylacyl, arylacyl, alkylsulfonyl, and arylsulfonyl. Preferred R2 is selected from hydrogen; C₁-C₆ alkyl, such alkyl being saturated or unsaturated with one double bond and unsubstituted or substituted with phenyl; C₁-C₆ alkylacyl, the alkyl being saturated or unsaturated with one double bond; and phenylacyl. More preferred is the alkyl portions of the aforementioned moieties being C₁-C₄ and saturated. More preferred still is R2 being methyl. Most preferred R2 is hydrogen.

In structure 1, each R3 is each independently selected from hydrogen, halo, alkyl, aryl, heterocycle, nitro and cyano; also from hydroxy, thio, amino, amide, formyl (acyl), carboxy, and carboxamide which are unsubstituted or substituted, preferably with alkyl or aryl or heterocycle; or two R3 together are alkylene or heteroalkylene attached to adjacent carbon atoms of the phenyl ring, thereby forming a cycloalkyl or aryl or heterocycle ring which is fused to the phenyl ring. Preferred R3 are independently selected from hydrogen, halo, alkyl, aryl, heterocycle, hydroxy, alkoxy, aryloxy, thio, alkylthio, arylthio, amino, alkylamino, arylamino, amide, alkylamide, arylamide, formyl, alkylacyl, arylacyl, carboxy and its alkyl or aryl esters and amides; more preferably from hydrogen, halo, C₁-C₄ alkyl, thio, C₁-C₄ alkylthio, C₁-C₄ mono- or dialkylamino, and C₁-C₄ alkylacyl. More preferred is for from one to three R3's being halo, the others being hydrogen. Also more preferred is for from one to three R3's being methyl or ethyl, the others being

hydrogen. Also preferred is one R3 being dialkylamino, the alkyls having from 1 to about 6 carbon atoms, preferably from about 1 to about 4 carbon atoms, and the others being hydrogen, such R3 preferably being attached to the 4' carbon. More preferred still is for from one to three R3's being independently selected from F, Cl and Br, the others being hydrogen; still more preferred, when two or three R3's are F, Cl or Br, they are the same. Also more preferred is from one to three R3's being unsubstituted methyl, the others being hydrogen. Also more preferred is one or two R3's being trifluoromethyl, the others being hydrogen.

Also preferred are two R3's which are attached to adjacent carbon atoms of the phenyl ring, together being a saturated or unsaturated alkylene or heteroalkylene having from 1 to about 6 carbon atoms and from 0 to about 3 heteroatoms, thus forming a ring fused to the phenyl, such ring having from about 5 to about 8 ring atoms. Such ring fused to the phenyl preferably has from about 5 to about 6 ring atoms of which from 0 to 2, more preferably 0 or 1, are heteroatoms. Preferred fused rings (including the phenyl to which the R3's are attached) include naphthyl, indolyl, benzimidazolyl, benzofuryl, benzopyranyl. When two R3's form a ring fused with the phenyl, other R3's are preferably H.

In structure 1, R4 is selected from hydrogen, halo, alkyl, aryl, heterocycle, carboxy and its alkyl esters and amides. Preferred R4 is selected from hydrogen, halo, C₁-C₄ alkyl, phenyl. More preferred R4 is selected from hydrogen and unsubstituted and substituted phenyl; substituents on such phenyl are preferably selected from hydroxy, alkoxy, thio and alkylthio. Most preferred R4 is hydrogen.

In structure 1, each R5 is independently selected from hydrogen, alkyl and aryl. Preferred R5 are selected from hydrogen and alkyl having from 1 to about 4 carbon atoms, especially unsubstituted methyl or ethyl. Most preferred is for both R5 to all be hydrogen.

In structure 1, each R6 is independently selected from hydrogen, halo, alkyl, aryl, heterocycle, cyano and nitro; also from hydroxy, thio, amino, amide, formyl (acyl), carboxy, carboxamide, and sulfonamide which are unsubstituted or substituted, preferably with alkyl or aryl or heterocycle. Preferred R6's are selected from hydrogen, halo, alkyl,

aryl, heterocycle, hydroxy, alkoxy, aryloxy, thio, alkylthio, arylthio, amino, alkylamino, arylamino, amide, alkylamide, arylamide, sulfonamide, alkylsulfonamide, arylsulfonamide, formyl, alkylacyl, arylacyl, carboxy and its alkyl and aryl esters and amides; more preferably from hydrogen, halo, hydroxy, C₁-C₄ alkoxy, thio, C₁-C₄ alkylthio, C₁-C₄ alkyl esters and amides of carboxy, and heterocycle having 5 or 6 ring atoms. More preferably one or both R₆'s are selected such that at least one heteroatom is bonded directly to the phenyl ring. When both R₆'s have heteroatoms bonded directly to the adjacent carbons of the phenyl ring, the two R₆'s may together form an alkylene moiety connecting the two heteroatoms, the alkylene moiety preferably having from 1 to about 4, more preferably 1 or 2, carbon atoms. Preferred is that one or both of the R₆'s are non-hydrogen moieties; more preferred is that both of them are non-hydrogen moieties. More preferred still is that both R₆'s are alkoxy or both alkylthio; preferably both are the same. Preferred alkyl portions of R₆'s have from 1 to about 4 carbon atoms, more preferably 1 or 2 carbon atoms; most preferred is methyl. Such alkyl portions are preferably unsubstituted. Such alkyl portions are preferably saturated. Most preferred is that both R₆'s are methoxy or ethoxy, especially methoxy.

Also preferred are a R₅ and a R₆, which are attached to adjacent carbon atoms of the phenyl ring, together being a saturated or unsaturated alkylene or heteroalkylene having from 1 to about 6 carbon atoms and from 0 to about 3 heteroatoms, thus forming a ring fused to the phenyl, such ring having from about 5 to about 8 ring atoms. Such ring fused to the phenyl preferably has from about 5 to about 6 ring atoms of which from 0 to 2, more preferably 0 or 1, are heteroatoms. Preferred rings formed by such R₅ and R₆ include phenyl, furyl, pyrrolyl, dioxanyl, imidazolyl, pyridinyl, pyrrolidinyl, piperidinyl. When such R₅ and R₆ form a fused ring, the other R₅ and R₆ are both preferably hydrogen.

In structure 1, B is nil or -C(R₇)=C(R₇)-. When B is nil, n is an integer from 0 to about 3, preferably 1 or 2. When B is -C(R₇)=C(R₇)-, n is an integer from 0 to about 1, preferably 0. Preferred B is nil.

In structure 1 and in B, each R₇ is independently selected from hydrogen, alkyl, and aryl. Non-hydrogen R₇ are preferably phenyl, or alkyl having from 1 to about 4


carbon atoms, preferably 1 or 2 carbon atoms. Such non-hydrogen R7 are preferably unsubstituted. Alkyl R7 are preferably saturated. Preferably no more than one of all the R7's is other than hydrogen. More preferably all R7's are hydrogen.

The subject invention includes optical isomers, diastereomers, and enantiomers of the compounds of structure 1. The subject invention includes pharmaceutically-acceptable salts, hydrates, and biohydrolyzable esters, amides and imides of such compounds.

A "pharmaceutically-acceptable salt" is a cationic salt formed at any acidic group (e.g., carboxy group), or an anionic salt formed at any basic group (e.g., amino group) on a compound of structure 1. Many pharmaceutically-acceptable salts are known. Preferred cationic salts include the alkali metal salts, such as sodium and potassium, alkaline earth metal salts, such as magnesium and calcium, and organic salts, such as ammonium. Preferred anionic salts include halides, sulfonates, carboxolates, phosphates, and the like. Salts of addition may provide an optical center where once there was none.

The compounds of the subject invention, and salts thereof, may have one or more chiral centers. The invention includes all optical isomers of the compounds of structure 1 and salts thereof, including diastereomers and enantiomers. The subject invention includes and contemplates each optical isomer, diastereomer or enantiomer thereof, in purified form, substantially purified form, and mixtures, including racemic mixtures.

Preferred compounds of the subject invention include those having structure 2:



In structure 2, R1's, R3's and R6's are as described hereinabove, and x is 0 or 1.

In structure 2, each R1 is preferably independently selected from linear alkanyl having from one to four carbon atoms, linear alkenyl having one double bond and from two to four carbon atoms, branched and cyclic alkanyl having from three to five carbon atoms, and branched and cyclic alkenyl having one double bond and from three to five

carbon atoms. Such preferred R1 are unsubstituted or substituted with one phenyl, more preferably are unsubstituted. More preferred R1 are selected from methyl, ethyl, ethenyl, n-propyl, i-propyl, n-propenyl, i-propenyl, s-butyl, cyclopropyl, cyclobutyl, and cyclopentyl. More preferred still R1 are selected from methyl, ethyl, ethenyl, i-propyl, and n-propenyl. Most preferred R1 are methyl. Also preferred is for both R1 to be the same.

In structure 2, one or both of R6, preferably both, are preferably alkylthio or more preferably alkoxy with alkanyl having from one to four carbon atoms. If one R6 is not alkylthio or alkoxy, it is preferably hydrogen. More preferred is both R6 being methoxy or ethoxy; most preferred is both R6 being methoxy.

In structure 2, the R3's are preferably selected from all hydrogen; mono-, di-, or trihalo, preferably selected from fluoro, chloro and bromo, preferably in one or more of the 2', 3', 4' and 5' positions; mono- di-, and trimethyl, preferably in one or more of the 2', 3', 4' and 6' positions; and mono- or di-trifluoromethyl, preferably in one or both of the 3' and 5' positions. Also preferred is one R3 being diakylamino, preferably in the 4' position, the two alkyls preferably being the same and preferably having from 1 to 4 carbon atoms, and the other R3's being hydrogen. More preferred combinations of R3's are selected from 4'-fluoro, 4'-chloro, 4'-bromo, 2',4'-difluoro, 2',4'-dichloro, 2',4'-dibromo, 2',4',5'-trifluoro, 2',4',5'-trichloro, 3',4'-difluoro, 3',4'-dichloro, 3',4'-dibromo, 4'-methyl, 2',4'-dimethyl, 2',4',6'-trimethyl, 3'-trifluoromethyl, 3',5'-di-trifluoromethyl, and 4'-dibutylamino; in each case all other R3's being hydrogen. Also preferred R3 combinations are selected from 2',4'-dihalo and 3',4'-dihalo, where one halo is selected from fluoro, chloro, and bromo, and the other halo is a different one of those three; more preferably one of such halo is fluoro, all other R3's having hydrogen. Most preferred R3 combinations are selected from 4'-chloro, 4'-bromo, and 2',4'-dichloro, all other R3's being hydrogen.

Non-limiting examples of compounds of the subject invention include those of structure 2 wherein both R6's are methoxy, and R1's and R3's are as indicated in the following table (R3's not specified are all hydrogen):

<u>Example</u>	<u>x</u>	<u>R1</u>	<u>R3</u>
----------------	----------	-----------	-----------

1	0	methyl, methyl	hydrogen
2	0	methyl, methyl	4'-fluoro
3	1	methyl, methyl	4'-chloro
4	0	methyl, methyl	4'-bromo
5	0	methyl, methyl	2',4'-difluoro
6	1	methyl, methyl	2',4'-dichloro
7	0	methyl, methyl	2'-fluoro, 4'-chloro
8	1	methyl, methyl	2'-fluoro, 4'-bromo

<u>Example</u>	<u>X</u>	<u>R1</u>	<u>R3</u>
9	0	methyl, methyl	2'-chloro, 4'-fluoro
10	1	methyl, methyl	2'-bromo, 4'-fluoro
11	1	methyl, methyl	3',4'-difluoro
12	0	methyl, methyl	3',4'-dichloro
13	0	methyl, methyl	3'-fluoro, 4'-chloro
14	1	methyl, methyl	3'-fluoro, 4'-bromo
15	0	methyl, methyl	3'-chloro, 4'-fluoro
16	1	methyl, methyl	3'-bromo, 4'-fluoro
17	0	methyl, methyl	2',4',5'-trifluoro
18	0	methyl, methyl	4'-methyl
19	1	methyl, methyl	2',4'-dimethyl
20	0	methyl, methyl	3',4'-dimethyl
21	1	methyl, methyl	2',4',6'-trimethyl
22	1	methyl, methyl	3'-trifluoromethyl
23	0	methyl, methyl	3',5'-di-trifluoromethyl
24	0	methyl, methyl	4'-fluoro, 3'-trifluoromethyl
25	1	ethyl, ethyl	4'-fluoro
26	0	ethyl, methyl	4'-chloro
27	0	ethyl, ethyl	4'-bromo
28	1	ethyl, methyl	2',4'-dichloro
29	1	ethyl, ethyl	3'-fluoro, 4'-chloro

30	1	ethenyl, ethenyl	4'-chloro
31	0	ethenyl, methyl	2',4'-dichloro
32	0	i-propyl, i-propyl	4'-fluoro
33	0	i-propyl, methyl	4'-chloro
34	1	i-propyl, i-propyl	4'-bromo
35	1	i-propyl, methyl	2',4'-dichloro
36	0	i-propyl, i-propyl	3'-fluoro, 4'-chloro
37	0	-CH ₂ -CH=CH ₂ , methyl	4'-fluoro
38	1	-CH ₂ -CH=CH ₂ , -CH ₂ -CH=CH ₂	4'-chloro

Example	x	R1	R3
39	1	-CH ₂ -CH=CH ₂ , ethyl	2',4'-dichloro
40	0	s-butyl, s-butyl	4'-chloro
41	1	cyclopentyl, methyl	4'-chloro
42	1	methyl, methyl	2',4',5'-trifluoro
43	0	methyl, methyl	2',3',4'-trifluoro
44	0	ethyl, ethyl	2',4',5'-trifluoro
45	1	ethyl, ethyl	2',3',4'-trifluoro
46	1	isopropyl, isopropyl	2',4',5'-trifluoro
47	0	isopropyl, isopropyl	2',3',4'-trifluoro
48	0	methyl, methyl	4'-dibutylamino
49	1	ethyl, ethyl	4'-dimethylamino
50	0	isopropyl, isopropyl	4'-diethylamino

In addition, it is recognized that for purification, administration, and the like, the salts and other derivatives of the above compounds can be used. Thus a pharmaceutically-acceptable salt, hydrate, or biohydrolyzable ester, amide or imide thereof is contemplated as part of the subject invention.

Methods of Making the Compounds


In making the compounds of the subject invention, the order of synthetic steps may be varied to increase yield of desired product. The skilled artisan will recognize that the judicious choice of reactants, solvents, and temperatures is important in successful synthesis. The starting materials used in preparing the subject compounds are known, made by known methods, or are commercially available.



It is recognized that the skilled artisan can readily carry out standard manipulations of organic compounds without further direction. These include, but are not limited to, reduction, oxidation, acylation, substitution, etherification, esterification, sulfonation, and the like. Examples of these manipulations are discussed in standard texts.

Procedures for preparing some imidazo-isoquinoline compounds are disclosed in U.S. Patent No. 3,917,610 issued November 4, 1975, and U.S. Patent No. 4,143,143 issued on March 6, 1979, both of which are incorporated herein by reference.


The following general schemes can be used for synthesizing compounds of the subject invention. In these schemes, R1, R2, R3, R4, R5, R6, R7, B, and n are as defined hereinabove. Non-conventional symbols and abbreviations for other moieties and chemicals, which may not be clear to the skilled chemist, are defined in the Examples hereinbelow.

General Scheme I








$\text{Pd(PPh}_3)_4$
Dioxane




$\text{MsCl/Et}_3\text{N}$
 CH_2Cl_2




$n\text{BuLi/THF}$




General Scheme II






The following examples provide further information regarding synthesis of the subject invention compounds.

Precursor Example 1




A suspended solution of NaH (6.5 g, 0.162 mol, washed with hexane twice) in anhydrous DMF (300 ml) is cooled in an ice/acetone bath (bath temp. -15 °C). Solid imidazole (10 g, 0.145 mol) is added in small portions and the mixture is stirred at room temperature (r.t.) for 0.5 h; the solution becomes clear. SEM-Cl (25 g, 0.150 mol) is added dropwise by a syringe pump at r.t. over 1 h; NaCl precipitates during the addition. The mixture is stirred at r.t. for about 1 h. Progress of the reaction is monitored by TLC (CH₂Cl₂/MeOH, 9:1). H₂O (10 ml) is added with caution to quench the reaction. The solvent is evaporated in vacuo. The residue is dissolved in Et₂O (200 ml) and washed with H₂O (4 x 50ml), brine (50 ml), dried (MgSO₄), filtered, and evaporated in vacuo to give compound A as an orange liquid.

Precursor Example 2




To a solution of SEM-protected imidazole A (1.48 g, 7.50 mmol) in dry THF (75 ml) under argon at -78 °C, n-BuLi (1.6 M in hexane) (6 ml, 9.60 mmol) is added dropwise and the mixture is stirred at -78 °C for 30 min. Phenyl disulfide (2.1 g, 9.60 mmol) in THF (2 ml) is then added dropwise. The dry ice/acetone bath is replaced with an ice bath after this addition. The mixture is stirred at 0 °C for 1 h, then at r.t. for 1 h. Progress is monitored by TLC (CH₂Cl₂/MeOH, 9:1). H₂O (5 ml) is added to quench the reaction. The solvent is evaporated in vacuo, and the residue is dissolved in Et₂O, washed with 5% NaHCO₃ (3 x 20ml), brine (20 ml), dried (MgSO₄), evaporated in vacuo, and purified by chromatography (silica gel, hexane/EtOAc 3:1) to give B as a yellow oil.

Precursor Example 3



3-chloroperoxybenzoic acid (MCPBA, 80-85%) (17.03 g, 78.9 mol) is added to a solution of SEM-protected 2-phenylsulfide imidazole **B** (9.71 g, 31.6 mmol) in anhydrous CH_2Cl_2 (160 mL), and the reaction is stirred under argon at room temperature for 15 hours. Progress monitored by TLC (hexane/EtOAc, 3:1). Sodium thiosulfate (3.9 g) is added to remove excess MCPBA. The mixture is filtered. The filtrate is washed with 5% Na_2CO_3 (3 x 150 mL), brine (150 mL), dried (MgSO_4), filtered, evaporated in vacuo, and purified by chromatography (silica gel, hexane/EtOAc 3:1) to give **C** as a light yellow oil.


Precursor Example 4



To a solution of SEM-protected 2-phenylsulfone imidazole **C** (8.61 g, 25.4 mmol) in anhydrous THF (250 mL) under argon at -78°C , $n\text{-BuLi}$ (1.6 M in hexane) (19.0 mL, 30.0 mmol) is added dropwise by a syringe pump; the solution is stirred at -78°C for 30 minutes. Tributyltin chloride (6.9 mL, 25.4 mmol) is added dropwise by a syringe pump. The mixture is stirred at room temperature for one hour. Progress is monitored by TLC (hexane/EtOAc, 9:1). H_2O (30 mL) is added to quench the reaction. The solvent is evaporated in vacuo. The residue is dissolved in ether (550 mL) and washed with saturated NH_4Cl (3 x 150 mL), brine (150 mL), dried (MgSO_4), filtered, evaporated in vacuo, and purified


by chromatography (silica gel, gradient: hexane (500 mL), hexane/EtOAc, 50:1; hexane/EtOAc, 12:1) to give **D** as a clear oil.

Precursor Example 5




$\text{Pd(PPh}_3)_4$ (0.0177 g, 0.015 mmol) is added to a solution of stannylimidazole **D** (0.51 g, 0.80 mmol), 4,5-dimethoxy-2-(2-hydroxyethyl)phenyl bromide **E** (0.33 g, 1.1 mmol), and LiCl (0.087 g, 2.1 mmol) in anhydrous dioxane (4.0 mL) at room temperature. A few crystals (~ 2 mg) of a radical scavenger, 2,6-di-*tert*-butyl-4-methylphenol, is added and the reaction is heated to reflux under argon for 5 hours. The reaction is cooled to room temperature and treated with a 1:1 mixture of ether and saturated aqueous KF solution (10 mL) for 15 hours. Progress is monitored by TLC (hexane/EtOAc, 3:1). The mixture is filtered through a pad of Celite with ether rinses. The filtrate is washed with water (3 x 12 mL), brine (3 x 12 mL), dried (MgSO_4), filtered, evaporated in vacuo, and purified by chromatography (silica gel, hexane/EtOAc, 2:3) to give **F** as an orange oil.

Precursor Example 6





To a solution of **E** (2.2 g, 4.24 mmol) and Et₃N (886 μ L) in dichloromethane (200 mL) under argon atmosphere at 0 °C, methylsulfonyl chloride (MsCl, 492 μ L) is added over a period of 0.5 h. The reaction is then warmed to room temperature and stirred for 1 h. TLC (EtOAc:Hexane, 1:1) is used to monitor the reaction; it indicates that MsO-ester formation and the SEM-cleavage followed by ring closure occur in one pot. The mixture is diluted with CH₂Cl₂ (25 mL), and washed with cold HCl aqueous (0.5N), NaHCO₃ aqueous, H₂O, brine and dried over MgSO₄. Filtration and evaporation of solvent gives a yellow solid. Purification using preparative HPLC (EtOAc:hexane, a gradient from 1:1 to 1:0) provides product **G** (1.3 g).

Invention Compound Example I




In a 25 mL single neck round bottom flask, equipped with a magnetic stir bar, argon inlet, and rubber septum, 1-methyl-1-phenyl-ethylamine (**H**) (164 mg, 1.2 mmol) in anhydrous THF under argon atmosphere is cooled to 0 °C. nBuLi in hexane (0.5 mL, 2.4 M) is added to the solution slowly. The reaction turns light yellow. After stirring for 45 minutes, compound **G** (150 mg, 0.405 mmol) in THF (1 mL) is added. The solution is warmed to room temperature, and is further heated to reflux for 12 h. TLC (CH₂Cl₂:CH₃OH, 99:1) indicates completion of the reaction. The solution is quenched with MeOH and evaporated to give a residue which is redissolved in CH₂Cl₂, washed with aqueous NaHCO₃ (5%), H₂O, brine, and dried over Na₂SO₄. The crude product, after filtration and evaporation in high vacuum in order to remove any excess amine **H**, is purified by chromatography (CH₂Cl₂:CH₃OH, 99:1) to provide subject invention compound **J**.

Invention Compound Example II




In a 25 mL single neck round bottom flask, compound J (181 mg, 0.5 mmol) in anhydrous THF (10 mL) under argon atmosphere is cooled to 0 °C, then NaH (18 mg) pre-washed with hexane is added as a suspension in hexane to the solution. After stirring for 15 minutes, methyl iodide (71 mg, 0.5 mmol) in THF (0.5 mL) is added. The solution is warmed to room temperature, and further heated to reflux for 2 h to complete the reaction. The solution is quenched with MeOH and evaporated to give a residue which is redissolved in CH₂Cl₂, washed with aqueous NaHCO₃ (5%), H₂O, brine and dried over Na₂SO₄. The crude product is purified by chromatography (CH₂Cl₂:CH₃OH from 99:1 to 95:5) to provide subject invention compound P.

Precursor Example 7




N-bromosuccinimide (NBS) solid (98 mg, 0.26 mmol) is added to a solution of compound **G** (0.5 mmol) in 15 mL of CCl_4 . Radical initiator benzoyl peroxide (2 mol%) is subsequently added. The flask is placed into a 90 °C oil bath. After 10 min stirring, the reaction is complete. Filtration of the mixture through a celite pad, and evaporation of the filtrate gives a residue. Purification by chromatography (EtOAc:hexane, 1:3 to 1:1) affords compound **K**.

Precursor Example 8



To a solution of **K** (0.22mmol) and Pd(PPh₃)₄ (13mg, 0.056mmol) in 7 mL of anhydrous toluene are added phenyltributyltin (0.26mmol) and a few crystals (~ 2 mg) of 2,6-di-*tert*-butyl-4-methylphenol. The reaction mixture is allowed to reflux at 110 °C under nitrogen for 6 hours to complete the reaction. The reaction mixture is allowed to cool, and is then diluted with 1-2 mL of ethyl acetate (EtOAc). The resultant mixture is washed with water, then brine, extracted with CH₂Cl₂, dried over Na₂SO₄, and filtered. The filtrate is treated with 3 mL of 30% aqueous KF at room temperature for 2h. The solid is filtered off. The filtrate is diluted with CH₂Cl₂ and washed with water, 30% aqueous NH₄OH (3X), brine, extracted with EtOAc, dried (Na₂SO₄), and concentrated *in vacuo* to yield crude product. Chromatography purification (silica gel, EtOAc:hexane, 1:1 to 1:0) yields compound **L**.

Subject Invention Example III



In a 25 mL single neck round bottom flask, equipped with a magnetic stir bar, argon inlet, and rubber septum, (1-methyl-1-phenyl-ethylamine) (**M**) (1.2 mmol) in anhydrous THF under argon atmosphere is cooled to 0 °C. *n*BuLi in hexane (0.5

mL, 2.4 M) is added slowly to the solution. The reaction turns light yellow. After stirring for 45 minutes, compound L (180 mg, 0.405 mmol) in THF (1 mL) is added. The solution is warmed to room temperature, and is further heated to reflux for 12 h to complete the reaction. The solution is quenched with MeOH and evaporated to give a residue which is redissolved in CH₂Cl₂, washed with aqueous NaHCO₃ (5%), H₂O, brine and dried over Na₂SO₄. The crude product, after filtration and evaporation in high vacuum in order to remove excess amine M, is purified by chromatography to provide subject invention compound N.

Compositions of the Invention

A composition of the subject invention comprises:

- a) a safe and effective amount of a compound of the invention; and
- b) a pharmaceutically-acceptable excipient.

Typically, such composition comprises several excipients. It may also optionally comprise other active compounds which do not substantially interfere with the activity of the subject invention compound.

The compositions of the subject invention may be in any of a variety of forms, suitable (for example) for oral, rectal, topical or parenteral administration. Compositions of the subject invention are preferably provided in unit dosage form. As used herein, a "unit dosage form" is a composition containing an amount of a subject compound that is suitable for administration to a human or lower animal subject, in a single dose, according to good medical practice.

As used herein, a "safe and effective amount" of a subject compound is an amount large enough to significantly induce a positive modification in the symptoms and/or condition to be treated in a host, but small enough to avoid serious adverse side effects in the host (such as toxicity, irritation, or allergic response), commensurate with a reasonable benefit/risk ratio. The safe and effective amount will vary with such factors as the particular condition being treated, the age and physical condition of the patient, the duration of treatment, the nature of concurrent therapy (if any), the specific dosage form to be used, and the dosage regimen employed.

The term "pharmaceutically-acceptable excipient", as used herein, includes physiologically inert, pharmacologically inactive substances which are compatible with the physical and chemical characteristics of the subject invention compound used, and which are of sufficiently high purity and sufficiently low toxicity to be suitable for administration to a human or lower animal. The term "compatible", as used herein, means that the excipients of the subject composition are capable of being commingled with the subject invention compound, and with each other in a manner such that there is no interaction which would substantially reduce the pharmaceutical efficacy of the compound, under ordinary use situations.

Depending upon the particular route of administration desired, a variety of pharmaceutically-acceptable excipients well-known in the art may be used. Excipients include, but are not limited to, polymers, resins, plasticizers, fillers, binders, lubricants, glidants, disintegrants, solvents, co-solvents, buffer systems, surfactants, preservatives, sweetening agents, flavoring agents, and dyes or pigments. The amount of excipients employed in conjunction with the subject compound is sufficient to provide a practical quantity of material for administration per unit dose of the subject compound.

Some examples of substances which can serve as pharmaceutically-acceptable excipients are sugars, such as lactose, dextrose, glucose and sucrose; starches, such as cornstarch and potato starch; cellulose and its derivatives, such as methylcellulose, sodium carboxymethylcellulose, ethylcellulose, hydroxypropylcellulose and cellulose acetate; polymers, such as povidone and carbomers; powdered tragacanth; gums, such as xanthan, guar and acacia; malt; solid lubricants, such as stearic acid, magnesium stearate, and talc; inorganic fillers, such as calcium phosphates and calcium sulfate; disintegrants, such as sodium starch glycolate, crospovidone, croscarmellose sodium, and microcrystalline cellulose; encapsulating and coating materials, such as gelatins, waxes, and cellulose derivatives; vegetable oils, such as peanut oil, cottonseed oil, sesame oil, olive oil, corn oil and oil of the obroma; polyols such as propylene glycol, glycerin, sorbitol, mannitol, and polyethylene glycol; alginic acid; surfactants such as the Tweens®, alkyl sulfate salts, salts of fatty acids, sucrose esters; ethyl oleate; coloring agents; flavoring agents; tableting agents; stabilizers; antioxidants; preservatives;

solvents, such as ethanol, pyrogen-free water; isotonic saline; and buffer solutions, such as phosphoric, tartaric, citric, and acetic acids, and their sodium, potassium, and ammonium salts.

Preferred compositions of the subject invention are oral dosage forms. The term "oral dosage form", as used herein, means any pharmaceutical composition intended to be systemically administered to an individual by delivering the composition via the mouth to the gastrointestinal tract of an individual. Preferred are oral unit dosage forms, such as tablets, coated or non-coated, and capsules, hard or soft gel. Subject oral unit dosage form compositions comprise preferably at least about 4 mg, more preferably at least about 20 mg, more preferably still at least about 100 mg, and preferably at most about 1000 mg, more preferably at most about 500 mg, more preferably still at most about 250 mg, of a subject compound. Subject oral dosage form compositions comprise preferably at least about 1%, more preferably at least about 10%, and preferably at most about 70%, more preferably at most about 40%, of a subject compound; and comprise preferably at least about 30%, more preferably at least about 60%, and preferably at most about 99%, more preferably at most about 90%, pharmaceutically-acceptable excipients.

Parenteral dosage forms are also preferred subject invention compositions. The term "parenteral dosage form", as used herein, means any pharmaceutical composition intended to be systemically administered to a human or lower animal via delivery of a solution or emulsion containing the active ingredient, by puncturing the skin of the individual, in order to deliver the solution or emulsion to the circulatory system of the individual either by intravenous, intramuscular, intraperitoneal or subcutaneous injection. Subject parenteral unit dosage form compositions comprise preferably at least about 1 mg, more preferably at least about 6 mg, more preferably still at least about 30 mg, and preferably at most about 400 mg, more preferably at most about 100 mg, more preferably still at most about 40 mg, of a subject compound. Subject parenteral dosage form compositions comprise preferably at least about 1%, more preferably at least about 5%, and preferably at most about 20%, more preferably at most about 10%, of a subject compound; and comprises preferably at least about 80%, more preferably at least about 90%, and preferably at most about 99%, more preferably at most about 95%,

pharmaceutically-acceptable excipients. In addition, dosages for injection may be prepared in dried or lyophilized form. Such forms can be reconstituted with water, saline solution, or a buffer solution, depending on the preparation of the dosage form. Such forms may be packaged as individual dosages or multiple dosages for easier handling. Where lyophilized or dried dosages are used, the reconstituted dosage form is preferably isotonic, and at a physiologically compatible pH, and comprises the subject compound and excipients in the amounts and percentages indicated previously in this paragraph.

Methods of Treatment Using the Compounds

Subject invention compounds have demonstrated pharmacological activity in processes known to be associated with one or more of cardiovascular activity, inflammatory mechanisms, oncology, and regulation of protein transport from cells. The subject invention includes methods of using the above compounds of the subject invention for therapeutic or preventative treatment of one or more of the following diseases or disorders: congestive heart failure, arrhythmia, hypotension, cardiac reperfusion injury, arteriosclerosis, restenosis, vascular tone, bacterial infection, cancer, Kaposi's sarcoma, psoriasis, migraine, nasal congestion, allergic responses, rheumatoid arthritis, and osteoporosis. Such methods comprise administering to a human or lower animal in need of such treatment or prevention a safe and effective amount of a subject invention compound.


For preferred oral administration of compounds and/or compositions of the subject invention, preferably at least about 0.1 mg/kg, more preferably at least about 0.5 mg/kg, more preferably still at least about 2 mg/kg, and preferably at most about 20 mg/kg, more preferably at most about 5 mg/kg, more preferably still at most about 2 mg/kg, of a subject compound is administered to a human or lower animal, preferably at least about 1 time, more preferably at least about 2 times, and preferably at most about 4 times, more preferably at most about 2 times, daily. Treatment duration using such oral daily dosages is dependent on the disease or disorder being treated; it is preferably at least about 1 day, more preferably at least about 3 days, more preferably still at least 7 days, and preferably at most about 5 years, more preferably at most about 60 days, more preferably still at most about 15 days.

For preferred parenteral administration of compounds and/or compositions of the subject invention, preferably at least about 0.04 mg/kg, more preferably at least about 0.2 mg/kg, more preferably still at least about 1 mg/kg, and preferably at most about 10 mg/kg, more preferably at most about 4 mg/kg, more preferably still at most about 1 mg/kg, of a subject compound is administered to a human or lower animal, preferably at least about 1 time, more preferably at least about 2 times, and preferably at most about 4 times, more preferably at most about 2 times, daily. Treatment duration using such parenteral daily dosages is dependent on the disease or disorder being treated; it is preferably at least about 1 day, more preferably at least about 3 days, more preferably still at least 7 days, and preferably at most about 60 days, more preferably at most about 20 days, more preferably still at most about 5 days.

While particular embodiments of the subject invention have been described, it will be obvious to those skilled in the arts that various changes and modifications of the subject invention can be made without departing from the spirit and scope of the invention. It is intended to cover, in the appended claims, all such modifications that are within the scope of this invention.

What is claimed is:

1. A compound having the structure:



wherein:

- (a) each R1 is independently selected from the group consisting of hydrogen, alkyl, aryl, and heterocycle; preferably each R1 is independently selected from the group consisting of hydrogen; linear, branched and cyclic alkanyl and alkenyl; and phenyl;
- (b) R2 is selected from the group consisting of hydrogen, alkyl, alkylacyl, arylacyl, alkylsulfonyl, and arylsulfonyl; preferably each R2 is selected from the group consisting of hydrogen; linear, branched and cyclic alkanyl and alkenyl; alkylacyl, and phenylacyl;
- (c) each R3 is independently selected from the group consisting of hydrogen, halo, alkyl, aryl, heterocycle, nitro, cyano, and unsubstituted or alkyl- or aryl- or heterocycle-substituted hydroxy, thio, amino, amide, formyl (acyl), carboxy, and carboxamide; or two R3's together are alkylene or heteroalkylene attached to adjacent carbons of the phenyl to which the R3's are attached; preferably each R3 is independently selected from the group consisting of hydrogen, halo, alkyl, aryl, heterocycle, hydroxy, alkoxy, aryloxy, thio, alkylthio, arylthio, amino, alkylamino, arylamino, amide, alkylamide, arylamide, formyl, alkylacyl, arylacyl, carboxy and its alkyl and aryl esters and amides; or preferably two R3's together are alkylene or heteroalkylene and, with the carbons to which they are attached, are cycloalkyl, aryl or heterocycle ring;

- (d) R4 is selected from the group consisting of hydrogen, halo, alkyl, aryl, heterocycle, and carboxy and its alkyl and aryl esters and amides;
- (e) each R5 is independently selected from the group consisting of hydrogen, alkyl, and aryl;
- (f) each R6 is independently selected from the group consisting of hydrogen, halo, alkyl, aryl, heterocycle, nitro, cyano, and unsubstituted or alkyl- or aryl- or heterocycle-substituted hydroxy, thio, amino, amide, sulfonamide, formyl, carboxy, and carboxamide; or the two R6's together are alkylene or heteroalkylene, or a R5 and a R6 which are attached to adjacent carbons together are alkylene or heteroalkylene; preferably each R6 is independently selected from the group consisting of hydrogen, halo, alkyl, aryl, heterocycle, hydroxy, alkoxy, aryloxy, thio, alkylthio, arylthio, amino, alkylamino, arylamino, amide, alkylamide, arylamide, sulfonamide, alkylsulfonamide, arylsulfonamide, formyl, alkylacyl, arylacyl, carboxy and its alkyl and aryl esters and amides; or preferably the two R6's together are alkylene or heteroalkylene and, with the carbons to which they are attached, are cycloalkyl, aryl or heterocycle ring;
- (g) B is nil or $-C(R7)=C(R7)-$;
- (h) if B is nil, n is an integer from 0 to about 3, otherwise n is from 0 to about 1;
- (j) each R7 is independently selected from the group consisting of hydrogen, alkyl and aryl; preferably each R7 is hydrogen or alkyl having from 1 to about 4 carbon atoms;

wherein preferably R2, R4 and both R5 are hydrogen;

and an optical isomer, diastereomer or enantiomer thereof; a pharmaceutically acceptable salt, hydrate, or biohydrolyzable ester, amide or imide thereof.

2. The compound of Claim 1 wherein:

- (a) each R1 is independently unsubstituted C₁-C₅ linear, or C₃-C₅ branched or cyclic, alkanyl; or unsubstituted C₂-C₅ linear, or C₃-C₅ branched or cyclic, alkenyl having one double bond;

- (b) R₂ is selected from the group consisting of hydrogen; unsubstituted C₁-C₅ linear, or C₃-C₅ branched or cyclic, alkanyl; or unsubstituted C₂-C₅ linear, or C₃-C₅ branched or cyclic, alkenyl having one double bond;
 - (c) each R₃ is independently selected from the group consisting of hydrogen, halo, C₁-C₄ alkyl, thio, C₁-C₄ alkylthio, C₁-C₄ mono- or dialkylamino, C₁-C₄ alkylacyl;
 - (d) R₄ is selected from the group consisting of hydrogen, halo, C₁-C₄ alkyl, and phenyl;
 - (e) both R₅'s are hydrogen;
 - (f) one R₆ is selected from the group consisting of alkoxy, alkylthio, monoalkylamino, dialkylamino, the alkyl portions thereof being saturated and unsubstituted and having from 1 to about 4 carbon atoms; the other R₆ is selected from the group consisting of hydrogen, alkoxy, alkylthio, monoalkylamino, dialkylamino, the alkyl portions thereof being saturated and unsubstituted and having from 1 to about 4 carbon atoms; or the alkyl portions of the two R₆'s are connected forming an alkylene moiety of from 1 to about 4 carbon atoms wherein preferably both R₆'s are C₁-C₄ alkoxy or C₁-C₄ alkylthio; and
 - (g) no more than one R₇ is other than hydrogen; preferably all R₇'s are hydrogen.
3. The compound of Claim 1 or 2 wherein both R₆'s are methoxy.
4. The compound of Claim 2 wherein B is nil, and n is 1 or 2.
5. The compound of Claim 2 wherein B is -C(R₇)=C(R₇)-, and n is 0.
6. The compound of Claim 1 or 2 wherein from one to three R₃'s are independently selected from F, Cl and Br, the others being hydrogen; or from one to three R₃'s are

unsubstituted methyl, the others being hydrogen; or one or two R3's are trifluoromethyl, the others being hydrogen.

7. The compound of Claim 2 wherein both R1's are methyl.
8. A composition comprising:
 - (a) a safe and effective amount of a compound of Claim 1 or 2; and
 - (b) a pharmaceutically-acceptable excipient.
9. The use of a compound of Claim 1 or 2 for the manufacture of a medicament for the treatment or prevention of nasal congestion, migraine, cardiac arrhythmia, cardiac reperfusion injury, congestive heart failure or hypotension in a human or lower animal subject in need thereof, the treatment or prevention comprising administering to the subject, a safe and effective amount of a compound of Claim 1 or 2.
10. The use of a compound of Claim 1 or 2 for the manufacture of a medicament for the treatment or prevention of a disease or disorder caused by cellular protein transport including, but not limited to, osteoporosis, rheumatoid arthritis, allergic responses, restenosis, vascular tone, cancer, Kaposi's sarcoma, psoriasis, and bacterial infections in a human or lower animal subject in need thereof, the treatment or prevention comprising administering to the subject, a safe and effective amount of a compound of Claim 1 or 2.

INTERNATIONAL SEARCH REPORT

onal Application No

PCT/US 00/13413

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D487/04 A61K31/415 //(C07D487/04,235:00,221:00),
(C07D487/04,235:00,223:00)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A01P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, PAJ, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	FR 2 190 458 A (CHINOIN GYOGYSZER ES VEGYESZET) - 1 February 1974 (1974-02-01) example 8 -/--	1,9,10

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Z" document member of the same patent family

Date of the actual completion of the international search

5 September 2000

Date of mailing of the international search report

19/09/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Frelon, D

INTERNATIONAL SEARCH REPORT

Int. Application No.

PCT/US 00/13413

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE CHEMABS 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; BOKCHARD, U. ET AL: "The positive inotropic, antiarrhythmic and (sodium-potassium ion)-dependent ATPase inhibitory effects of the isoquinoline derivative, BIIA" retrieved from STN Database accession no. 94:41256 XP002146553 RN 52210-64-7 and marker & NAUNYN-SCHMIEDEBERG'S ARCH. PHARMACOL. (1980), 312(2), 187-92 ,</p>	1,9,10
X	<p>DATABASE CHEMABS 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; HALA, O. ET AL: "Electrical and mechanical activity of the myocardium as affected by CH-101, an antiarrhythmic agent with preferential supraventricular action" retrieved from STN Database accession no. 107:510 XP002146554 RN 90936-98-4 and markers & ADV. PHARMACOL. RES. PRACT., PROC. CONGR. HUNG. PHARMACOL. SOC., 4TH (1986), MEETING DATE 1985, VOLUME 1, 163-7. EDITOR(S): KNOLL, JOZSEF; KELEMEN, KAROLY. PUBLISHER: PERGAMON, OXFORD, UK. ,</p>	1,9,10
A	<p>FR 2 366 289 A (SANDOZ SA) 28 April 1978 (1978-04-28) page 7 -page 8; claims</p>	1-10
A	<p>PATENT ABSTRACTS OF JAPAN vol. 1997, no. 03, 31 March 1997 (1997-03-31) -& JP 08 291173 A (LEDERLE JAPAN LTD), 5 November 1996 (1996-11-05) abstract</p>	1-10
A,P	<p>WO 00 21964 A (PROCTER & GAMBLE) 20 April 2000 (2000-04-20) the whole document</p>	1-10
A,P	<p>WO 00 21961 A (PROCTER & GAMBLE) 20 April 2000 (2000-04-20) the whole document</p>	1-10

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/US 00/13413

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
FR 2190458	A	01-02-1974	HU 167240 B	27-09-1975
			AR 208055 A	30-11-1976
			AR 210066 A	30-06-1977
			AR 209330 A	15-04-1977
			AR 209331 A	15-04-1977
			AR 211857 A	31-03-1978
			AT 346333 B	10-11-1978
			AT 316077 A	15-03-1978
			AT 329058 B	26-04-1976
			AT 563773 A	15-07-1975
			AT 343663 B	12-06-1978
			AT 627175 A	15-10-1977
			AT 347465 B	27-12-1978
			AT 627275 A	15-05-1978
			AT 343664 B	12-06-1978
			AT 627375 A	15-10-1977
			BE 801668 A	15-10-1973
			CA 1014559 A	26-07-1977
			CH 602729 A	31-07-1978
			CH 602730 A	31-07-1978
			CH 602731 A	31-07-1978
			CH 603639 A	31-08-1978
			CH 603647 A	31-08-1978
			CH 610900 A	15-05-1979
			CS 179024 B	31-10-1977
			DD 108090 A	05-09-1974
			DE 2332860 A	07-03-1974
			DK 141066 B	07-01-1980
			DK 141067 B	07-01-1980
			DK 545577 A,B,	08-12-1977
			DK 545677 A,B,	08-12-1977
			DK 545777 A,B,	08-12-1977
			EG 11302 A	30-06-1978
			ES 416971 A	16-02-1976
			FI 55199 B	28-02-1979
			FI 772708 A,B,	14-09-1977
			FI 772709 A,B,	14-09-1977
			FI 772710 A,B,	14-09-1977
			FI 772711 A,B,	14-09-1977
			GB 1438819 A	09-06-1976
			IL 42613 A	31-05-1977
			IN 144283 A	22-04-1978
			IN 144284 A	22-04-1978
			IN 144285 A	22-04-1978
			IN 144287 A	22-04-1978
			IN 139710 A	24-07-1976
			JP 954929 C	31-05-1979
			JP 49042697 A	22-04-1974
			JP 53036480 B	03-10-1978
			JP 1150118 C	14-06-1983
FR 2366289	A	28-04-1978	CH 603643 A	31-08-1978
			AT 688677 A	15-11-1981
			AU 514995 B	12-03-1981
			AU 2920677 A	05-04-1979
			BE 859084 A	28-03-1978
			CA 1090804 A	02-12-1980
			DE 2742433 A	30-03-1978

INTERNATIONAL SEARCH REPORT

information on patent family members



Int'l. Application No

PCT/US 00/13413

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
FR 2366289 A		DK 416077 A,B, ES 462672 A FI 772758 A GB 1589752 A JP 53044598 A NL 7710414 A NZ 185275 A PH 12671 A PT 67084 A,B SE 7710504 A SU 639450 A US 4143143 A ZA 7705818 A	30-03-1978 01-12-1978 30-03-1978 20-05-1981 21-04-1978 31-03-1978 26-08-1980 11-07-1979 01-10-1977 30-03-1978 25-12-1978 06-03-1979 30-05-1979
JP 08291173 A	05-11-1996	NONE	
WO 0021964 A	20-04-2000	AU 6414599 A	01-05-2000
WO 0021961 A	20-04-2000	AU 6289399 A	01-05-2000



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁷ : C07D 471/04, 487/04, C07F 7/10 // A61K 31/5517, 31/437, A61P 9/00, 29/00, (C07D 471/04, 235:00, 221:00) (C07D 487/04, 243:00, 235:00)</p>	A1	<p>(11) International Publication Number: WO 00/69857</p> <p>(43) International Publication Date: 23 November 2000 (23.11.00)</p>
<p>(21) International Application Number: PCT/US00/13417</p> <p>(22) International Filing Date: 16 May 2000 (16.05.00)</p> <p>(30) Priority Data: 60/134,829 19 May 1999 (19.05.99) US</p> <p>(71) Applicant (for all designated States except US): THE PROCTER & GAMBLE COMPANY [US/US]; One Procter & Gamble Plaza, Cincinnati, OH 45202 (US).</p> <p>(72) Inventors; and</p> <p>(75) Inventors/Applicants (for US only): LIU, Song [CN/US]; 9338 Ambleside Drive, Cincinnati, OH 45241 (US). LI, Min [US/US]; 7563 Lakota Springs Drive, West Chester, OH 45069 (US).</p>		<p>(81) Designated States: AE, AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), DM, EE, EE (Utility model), ES, FI, FI (Utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published</p> <p>With international search report.</p>
<p>(74) Agents: REED, T., David et al.; The Procter & Gamble Company, 5299 Spring Grove Avenue, Cincinnati, OH 45217-1087 (US).</p>		<p>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</p>
<p>(54) Title: PROCESS FOR MAKING FUSED-RING IMIDAZO-CONTAINING COMPOUNDS</p>		
<div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;">  <p>(1)</p> </div> <div style="text-align: center;">  <p>(2)</p> </div> </div>		
<p>(57) Abstract</p> <p>The subject invention involves processes for making fused-ring imidazo-containing compounds having structure (1) by obtaining and cyclizing compounds having structure (2) wherein m, n, R, R1, Q1 and Q2 are as defined herein, -X1 is selected from -Cl, -Br, -I, -OH and -COOH, and -SEM is -CH2OCH2CH2-Si(CH3)3.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						